

CODIMENSION AND CONNECTEDNESS OF DEGENERACY LOCI OVER LOCAL RINGS

HUBERT FLENNER AND BERND ULRICH

ABSTRACT. We deduce results on the dimension and connectedness of degeneracy loci of maps of finite modules $f : M \rightarrow N$ over a local noetherian ring $(A, \mathfrak{m})$. We show for instance that the expected determinantal bounds on the dimension of the t -th degeneracy locus of f hold if $f \in \mathfrak{m}\mathrm{Hom}(M, N)$, and that this degeneracy locus is connected in the expected dimension provided $\hat{A}$ is a domain.

1. INTRODUCTION

In this paper we will investigate the dimension and connectedness properties of degeneracy loci of homomorphisms $f : M \rightarrow N$ of finite modules over a noetherian local ring $(A, \mathfrak{m})$. For $t \geq 0$ we consider the $(t+1)$ -th determinantal ideal $I_{t+1}(f)$ of f , which for free modules is just the ideal of $(t+1)$ -minors of a defining matrix of f (see 4.1 for the general definition). We will present two types of results for these ideals or, equivalently, the degeneracy loci $D_t(f) := \mathrm{Spec}(A/I_{t+1}(f))$, namely on

- (I) the dimension of $D_t(f)$,
- (II) the connectedness of $D_t(f)$.

In commutative algebra there are various results known in this direction. Concerning (I), the classical Krull principal ideal theorem states that the dimension drops at most by one when we factor out an element $x \in \mathfrak{m}$. More generally for determinantal ideals there is the well known Eagon-Northcott bound on the height of such ideals [EN].

The principal ideal theorem was generalized to order ideals by Eisenbud and Evans [EE] and in its general form by Bruns [Br]. More precisely, if N is a finite A -module, $f \in \mathfrak{m}N$ and $N^\vee(f) = \{\varphi(f) | \varphi \in \mathrm{Hom}_A(N, A)\}$ is the order ideal of f , then under suitable hypotheses the dimension of the quotient $A/N^\vee(f)$ is at least $\dim A - \mathrm{rk} N$. In the setup above this is the case where $M = A$ and $t = 0$, i.e. $N^\vee(f) = I_1(f)$. In the meantime there has been a wide range of extensions and applications of these results, see e.g. [EHU₁, EHU₃, EHU₄].

Concerning (II), much less is known. Using methods from formal geometry Grothendieck showed in [SGA, Exp. XIII] the analogue of the principal ideal theorem for connectedness. Grothendieck's theorem says in its simplest form that for a noetherian complete local domain A of dimension ≥ 3 and $x \in \mathfrak{m}$ the punctured spectrum $\mathrm{Spec}(A/xA) \setminus \{\mathfrak{m}\}$ is connected (see [FOV], Section 3.1 for a systematic treatment of connectedness and a more elementary proof of this result). In a more

This paper originated during visits of the authors at their home universities. They are grateful for the hospitality and the financial support by the DFG Schwerpunkt "Globale Methoden der komplexen Analysis". The second author was supported in part by the NSF.

recent paper F. Steffen [St] generalized this to determinantal ideals of matrices; however her method is restricted to characteristic 0.

Our main results are as follows. First we show: *if $f \in \mathfrak{m} \operatorname{Hom}_A(M, N)$ and if for some minimal prime $\mathfrak{q}$ of A of maximal dimension the modules $M_{\mathfrak{q}}$ and $N_{\mathfrak{q}}$ are free over $A_{\mathfrak{q}}$ of ranks m and n , respectively, then $\dim D_t(f) \geq \dim A - \tau$ with $\tau := (n - t)(m - t)$. This generalizes the Bruns-Eisenbud-Evans principal ideal theorem for order ideals. If moreover M and N are free at all primes $\neq \mathfrak{m}$ and $\hat{A}$ is connected in dimension d , then $D_t(f)$ is connected in dimension $d - \tau$. Applying this to free modules we recover the connectedness theorem of F. Steffen mentioned above in any characteristic. Recall that a noetherian scheme Y is said to be *connected in dimension d* for d an integer, if $\dim Y > d$ and if $Y \setminus X$ is connected for every closed subset $X \subseteq Y$ with $\dim X < d$.*

In projective geometry Fulton and Lazarsfeld showed analogous results for maps of vector bundles $f : \mathcal{M} \rightarrow \mathcal{N}$ if $\mathcal{H}om(\mathcal{M}, \mathcal{N})$ is ample. Ideally the local statements should imply the corresponding projective results by passing to the affine cones. However, the assumption that $f \in \mathfrak{m} \operatorname{Hom}(M, N)$ turns out to be too restrictive for such applications. We are able to obtain such results in characteristic $p > 0$ provided that $\mathcal{H}om(\mathcal{M}, \mathcal{N})$ is cohomologically p -ample in the sense of Kleiman [Kl]. To deduce this from a local statement we introduce in Section 2 as a new tool the concepts of (cohomological) p -ampleness and ampleness for modules over local rings that reflect to a certain extent the corresponding notions in projective geometry, see Propositions 2.4 and 2.8. This allows us to weaken the assumption “ $f \in \mathfrak{m} \operatorname{Hom}(M, N)$ ” to the condition that the pair $(Af, \operatorname{Hom}(M, N))$ is p -ample, see Theorems 4.2 and 4.4.

The A -dual of an A -module M will be denoted by $M^\vee$. If $(A, \mathfrak{m})$ is local we will say that M is a module of *rank m* with an *isolated singularity* if $M_{\mathfrak{p}}$ is free of constant rank m for all non-maximal primes $\mathfrak{p}$ of A . As a general reference for basic facts on commutative algebra and algebraic geometry we refer to the books of Eisenbud [Ei], Matsumura [Ma] and Hartshorne [Ha₂].

2. p -AMPLENESS AND AMPLENESS OVER LOCAL RINGS

In this section we propose a definition of p -ampleness and ampleness over local rings. Such notions should reflect the usual p -ampleness and ampleness in global geometry as introduced by Hartshorne [Ha₁]. We recall shortly the relevant definitions in the projective case.

Let $R = R_0[R_1]$ be a finitely generated standard graded algebra over a field $K = R_0$ and M a finitely generated graded R module such that the associated sheaf $\mathcal{M} = \tilde{M}$ on $X := \operatorname{Proj} R$ is locally free. Then $\mathcal{M}$ is called *ample* if for every other coherent sheaf $\mathcal{N}$ on X the sheaf $\mathbf{S}^n(\mathcal{M}) \otimes_X \mathcal{N}$ is globally generated for $n \gg 0$, where $\mathbf{S}^n$ denotes the n th symmetric power functor. We note that this definition is equivalent to the cohomological condition

$$(\#) \quad H^j(X, \mathbf{S}^n(\mathcal{M}) \otimes_X \mathcal{N}) = 0 \quad \text{for all } j > 0, n \geq n_0 = n_0(\mathcal{N})$$

and all $\mathcal{N}$ as above, see [Ha₁, 3.3].

If K has characteristic $p > 0$, p -*ampleness* means by definition that for every coherent sheaf $\mathcal{N}$ on X the sheaf $F^{a*}(\mathcal{M}) \otimes_X \mathcal{N}$ is globally generated for $a \gg 0$, where $F : X \rightarrow X$ is the Frobenius map, see [Ha₁, Sect. 6]. We note that one cannot characterize this property by a cohomological condition similar to the one in (#).

Thus S. Kleiman [Kl] introduced the notion of *cohomologically p -ample* locally free sheaves meaning that

$$H^j(X, F^{a*}(\mathcal{M}) \otimes_A \mathcal{N}) = 0 \quad \text{for all } j > 0, \ n \geq n_0 = n_0(\mathcal{N})$$

and all $\mathcal{N}$ as above.

The following implications are known:

$$\text{cohomologically } p\text{-ample} \xrightarrow{[\text{Kl}]} p\text{-ample} \xrightarrow{[\text{Ha}_1, 6.3]} \text{ample}.$$

However, the converse implications do not hold as shown by Kleiman [Kl] and Gieseker [Gi, p. 111].

These notions heavily depend on the grading. Thus to define a local analogue our idea is to consider a relative notion depending on a pair (M', M) , where M' is a submodule of M .

2.1. Let $(A, \mathfrak{m})$ be a noetherian local ring of prime characteristic $p > 0$ and M an A -module. The Frobenius map $F : A \rightarrow A' := A$ is a ring homomorphism, and

$$M \mapsto F(M) := M \otimes_F A = M \otimes_A A'$$

is a functor on the category of A -modules. Iterating a times we obtain the map $F^a : A \rightarrow A' := A$ and the corresponding functor F^a . In order to distinguish between duals into A and A' we write $M^\vee = \text{Hom}_A(M, A)$ and $M^\times = \text{Hom}_{A'}(M, A')$.

For our purposes the following definition of p -ampleness is the most appropriate:

Definition 2.2. A pair (M', M) , where $M' \subseteq M$ are A -modules, is called *p -ample* if for $a \gg 0$ the image of the natural map

$$F^a(M') \rightarrow F^a(M)^{\times \times}$$

is contained in $\mathfrak{m}F^a(M)^{\times \times}$. Similarly, for $R = R_0[R_1]$ a noetherian graded algebra over a local ring $A = R_0$ and $M' \subseteq M$ graded R -modules we say that the pair (M', M) is p -ample, if the condition above is satisfied over R with $\mathfrak{m}$ the maximal homogeneous ideal of R or, equivalently, if the pair $(M'_\mathfrak{m}, M_\mathfrak{m})$ is p -ample over the localized ring $A = R_\mathfrak{m}$.

We first record some basic properties of this notion:

Remark 2.3. (1) Any pair (M', M) with $M' \subseteq \mathfrak{m}M$ is p -ample.

(2) If the pair (M', M) is p -ample, then so is (N, M) for every submodule N of M' .

(3) A pair (M', M) of finite A -modules is p -ample if and only if the pair of $\hat{A}$ -modules $(\hat{M}', \hat{M})$ is p -ample.

(4) Let $f : M_1 \rightarrow M_2$ be a homomorphism of A -modules. If (M', M_1) is p -ample, then so is $(f(M'), M_2)$.

(5) If (M', M) is p -ample, then $(N \otimes_A M', N \otimes_A M)$ is p -ample for any A -module N .

(6) If (M', M) is p -ample, then so is $(M' \cdot \mathbf{S}^{n-1}M, \mathbf{S}^n M)$ for every $n > 0$.

Proof. Parts (1), (2), (3) and (4) are immediate from the definitions. The assertions (5) and (6) are an easy consequence of the fact that the Frobenius functor is compatible with tensor products and symmetric powers. $\square$

In the following proposition we describe the relationship of our notion of p -ampleness with the corresponding projective notion.

Proposition 2.4. *Let $R = R_0[R_1]$ be a finitely generated graded algebra over a field $K = R_0$ of characteristic $p > 0$ with $\text{depth } R \geq 2$ and let $M = \bigoplus_{i \in \mathbb{Z}} M_i$ be a finite graded R -module with an isolated singularity. Then the following hold:*

- (1) *If the associated locally free sheaf $\mathcal{M} := \tilde{M}$ on $X := \text{Proj } R$ is cohomologically p -ample, then the pair $(R \cdot M_0, M)$ is p -ample.*
- (2) *If $\mathcal{M}$ is generated by the sections in M_0 and if the pair $(R \cdot M_0, M)$ is p -ample, then $\mathcal{M}$ is p -ample on X .*

Proof. Since $\text{depth } A \geq 2$ and M has an isolated singularity we have

$$(*) \quad F^a(M)^{\times \times} \cong H^\bullet_\bullet(F^{a*}(\mathcal{M})) := \bigoplus_{i \in \mathbb{Z}} H^0(X, F^{a*}(\mathcal{M})(i)).$$

In order to show (1) let $x_1, \dots, x_k \in R_1$ be a K -basis. The Koszul complex $K(\underline{x}, F^a(M))$ on $\underline{x}$ provides an exact sequence

$$0 \longrightarrow F^{a*}(\mathcal{M})(-k) \longrightarrow \dots \longrightarrow F^{a*}(\mathcal{M})(-1)^{\binom{k}{1}} \longrightarrow F^{a*}(\mathcal{M}) \longrightarrow 0$$

on X . As $\mathcal{M}$ is cohomologically p -ample the groups $H^j(X, F^{a*}(\mathcal{M})(-i))$ vanish for $j > 0$ and $0 \leq i \leq k$, if $a \gg 0$. Thus taking sections the sequence

$$\dots \longrightarrow H^0(X, F^{a*}(\mathcal{M})(-1))^{\binom{k}{1}} \longrightarrow H^0(X, F^{a*}(\mathcal{M})) \longrightarrow 0$$

is exact. In view of $(*)$ this implies that $R_1 \cdot F^a(M)^{\times \times}_{-1} = F^a(M)^{\times \times}_0 \supseteq F^a(RM_0)_0$ as desired.

If the conditions in (2) are satisfied then

$$F^a(RM_0)_0 \subseteq R_1 F^a(M)^{\times \times}_{-1} = R_1 H^0(X, F^{a*}(\mathcal{M})(-1))$$

for $a \gg 0$. As $\mathcal{M}$ is globally generated by M_0 it follows that the map

$$\mathcal{O}_X(1) \otimes H^0(X, F^{a*}(\mathcal{M})(-1)) \longrightarrow F^{a*}(\mathcal{M})$$

is surjective. Hence $F^{a*}(\mathcal{M})$ is a quotient of a direct sum of copies of $\mathcal{O}_X(1)$, and $\mathcal{M}$ is p -ample as required. $\square$

Although not needed in the sequel it is interesting to note that one can also introduce a local notion of ampleness. For the remainder of this section let $(A, \mathfrak{m})$ be an arbitrary noetherian local ring.

Definition 2.5. A pair (M', M) , where $M' \subseteq M$ are A -modules, is called *ample* if for $n \gg 0$ the image of the natural map $\mathbf{S}^n M' \rightarrow (\mathbf{S}^n M)^{\vee \vee}$ is contained in $\mathfrak{m}(\mathbf{S}^n M)^{\vee \vee}$.

As in Definition 2.2 one can define a similar notion in the graded case. In analogy to 2.3 we note some basic properties of this notion.

Remark 2.6. (1) Any pair (M', M) with $M' \subseteq \mathfrak{m}M$ is ample.

(2) If the pair (M', M) is ample, then so is (N, M) for every submodule N of M' .

(3) A pair (M', M) of finite A -modules is ample if and only if the pair of $\hat{A}$ -modules $(\hat{M}', \hat{M})$ is ample.

(4) Let $f : M_1 \rightarrow M_2$ be a homomorphism of A -modules. If (M', M_1) is ample then so is $(f(M'), M_2)$.

(5) If $\mathbb{Q} \subseteq A$ and (M', M) is ample, then $(M' \otimes_A N, M \otimes_A N)$ is ample for any A -module N .

(6) If $\mathbb{Q} \subseteq A$ and (M', M) is ample, then so is $(M' \cdot \mathbf{S}^{n-1} M, \mathbf{S}^n M)$ for every $n > 0$.

Proof. The proofs of (1), (2), (3) and (4) are immediate from the definitions. Part (5) follows from the fact that in characteristic 0 there is a natural surjective map $N^{\otimes n} \otimes_A \mathbf{S}^n(M) \rightarrow \mathbf{S}^n(N \otimes_A M)$. Finally, part (6) is an easy consequence of (5) and (4), as the pair $(M' \otimes_A \mathbf{S}^{n-1}M, M \otimes_A \mathbf{S}^{n-1}M)$ is ample and maps onto the pair $(M' \cdot \mathbf{S}^{n-1}M, \mathbf{S}^n M)$. We leave the simple details to the reader. $\square$

Remark 2.7. By a well-known result of Barton [Ba] the tensor product of ample vector bundles on a projective variety is again ample in any characteristic. Therefore one should expect that 2.6(5),(6) remain valid in positive characteristic as well.

The following proposition explains our terminology of ample pairs. Its proof follows along the same line of arguments as in 2.4 and is left to the reader.

Proposition 2.8. *Let $R = R_0[R_1]$ be a finitely generated graded algebra over a field $K = R_0$ with $\text{depth } R \geq 2$ and let $M = \bigoplus_{i \in \mathbb{Z}} M_i$ be a finite graded R -module with an isolated singularity. Then the following hold :*

- (1) *If the associated locally free sheaf $\mathcal{M} := \tilde{M}$ on $X := \text{Proj } R$ is ample, then the pair $(R \cdot M_0, M)$ is ample.*
- (2) *If $\mathcal{M}$ is generated by the sections in M_0 and if the pair $(R \cdot M_0, M)$ is ample then $\mathcal{M}$ is ample.*

Let A be a noetherian local ring, let M be a finitely generated A -module, and consider an embedding of $M^{\vee\vee}$ into a free A -module F . We recall that the *integral closure* of a submodule $N \subseteq M$ is the set of all elements in M whose image in F is integral over the image of the symmetric algebra $\mathbf{S}(N)$ in $\mathbf{S}(F)$ (this definition does not depend on the chosen embedding $M^{\vee\vee} \subseteq F$, see [EHU₂]). We have the following relation with our notion of ampleness:

Proposition 2.9. *Let $(A, \mathfrak{m})$ be a noetherian local ring and let $M' \subseteq M$ be finite A -modules. Assume that $M_{\mathfrak{q}}$ is free for all associated primes $\mathfrak{q}$ of A . If $M' \subseteq \overline{\mathfrak{m}M}$ is in the integral closure of $\mathfrak{m}M$ in M , then the pair (M', M) is ample*

Proof. Write M^n for the image of $\mathbf{S}^n M$ in $\mathbf{S}^n F$. As M is free at the associated primes of A we have $(\mathbf{S}^n M)^{\vee\vee} = (M^n)^{\vee\vee}$. By our assumption, the image of $\mathbf{S}^n M'$ in M^n is contained in $\mathfrak{m}M^n$ for $n \gg 0$, and hence is contained in $\mathfrak{m}(M^n)^{\vee\vee} = \mathfrak{m}(\mathbf{S}^n M)^{\vee\vee}$. $\square$

3. MAIN TECHNICAL RESULTS

3.1. Let A be a ring, E an A -module and $f \in E^\vee$. The linear form $f : E \rightarrow A$ extends uniquely to a morphism of A -algebras denoted

$$\varphi_f : \mathbf{S}(E) \longrightarrow A,$$

where $\mathbf{S}(E) = \bigoplus_{i=0}^{\infty} (\mathbf{S}^i E) T^i$ is the symmetric algebra of E over A . Note that if $E_{\mathfrak{p}}$ is free for some prime $\mathfrak{p} \in \text{Spec } A$, then the localization of the symmetric algebra $\mathbf{S}_{A_{\mathfrak{p}}}(E_{\mathfrak{p}})$ is isomorphic to a polynomial ring over $A_{\mathfrak{p}}$ in $\text{rk}(E_{\mathfrak{p}})$ indeterminates.

The next theorem is one of our main technical results:

Theorem 3.2. *Let $(A, \mathfrak{m})$ be a noetherian local ring, E a finite A -module and $f \in E^\vee$. Assume that*

$$(*) \quad h \circ f \in \mathfrak{m}' \text{Hom}_A(E, A')$$

for some local homomorphism $h : A \rightarrow A'$ with $(A', \mathfrak{m}')$ a noetherian local ring and $\dim A'/\mathfrak{m}A' = 0$. Further suppose that there exists a prime ideal $\mathfrak{q}'$ of A' with contraction $\mathfrak{q} := h^{-1}(\mathfrak{q}')$ so that $\dim A'/\mathfrak{q}' = \dim A$ and $E_{\mathfrak{q}}$ is a free $A_{\mathfrak{q}}$ -module of rank r . Let $I \subseteq S := \mathbf{S}(E)$ be a homogeneous ideal with $I \cap A \subseteq \mathfrak{q}$ and $\dim S_{\mathfrak{q}}/I_{\mathfrak{q}} = r - \tau$. If $I(f) := \varphi_f(I)$ then $\dim A/I(f) \geq \dim A - \tau$.

We will reduce Theorem 3.2 to the special case where h is the identity map:

Theorem 3.3. *Let $(A, \mathfrak{m})$ be a noetherian local ring, E a finite A -module and assume that*

$$(**) \quad f \in \mathfrak{m}E^{\vee}.$$

Further suppose there exists a prime ideal $\mathfrak{q}$ of A so that $\dim A/\mathfrak{q} = \dim A$ and $E_{\mathfrak{q}}$ is a free $A_{\mathfrak{q}}$ -module of rank r . Let $I \subseteq S := \mathbf{S}(E)$ be a homogeneous ideal with $I \cap A \subseteq \mathfrak{q}$ and $\dim S_{\mathfrak{q}}/I_{\mathfrak{q}} = r - \tau$. If $I(f) := \varphi_f(I)$ then $\dim A/I(f) \geq \dim A - \tau$.

Proof of Theorem 3.2. We show how Theorem 3.2 can be deduced from Theorem 3.3. For this we consider $E' := E \otimes_A A'$, $f' := f \otimes 1 \in E'^{\vee}$, $S' := S \otimes_A A' = \mathbf{S}(E')$, and the extended ideal $I' := IS'$. Notice that $I'(f') = I(f)A'$. If $t_1, \dots, t_n \in \mathfrak{m}$ is a system of parameters of A , then $\dim A'/(t_1, \dots, t_n)A' = \dim A'/\mathfrak{m}A' = 0$ and so by Krull's principal ideal theorem $\dim A' \leq n = \dim A$. Likewise we have $\dim A'/I'(f') \leq \dim A/I(f)$. Since $\dim A'/\mathfrak{q}' \leq \dim A' \leq \dim A$ and the outer terms are equal, it follows that $\mathfrak{q}'$ is a minimal prime of A' having maximal dimension and that $\dim A' = \dim A$. As $S'_{\mathfrak{q}'}$ and $S_{\mathfrak{q}}$ are polynomial rings in the same r variables over the Artinian local rings $A'_{\mathfrak{q}'}$ and $A_{\mathfrak{q}}$, reducing modulo the nilradicals of the latter rings one sees that $\dim S'_{\mathfrak{q}'}/I'_{\mathfrak{q}'} = \dim S_{\mathfrak{q}}/I_{\mathfrak{q}} = r - \tau$. Thus Theorem 3.3 gives $\dim A'/I'(f') \geq \dim A' - \tau$. Therefore

$$\dim A/I(f) \geq \dim A'/I'(f') \geq \dim A' - \tau = \dim A - \tau,$$

proving Theorem 3.2. $\square$

It remains to prove Theorem 3.3. For this we will use the following setting:

3.4. Fix a set of generators $f_1, \dots, f_k$ for $E^{\vee}$ and let

$$\varphi : S = \mathbf{S}(E) \rightarrow A[\underline{Y}] := A[Y_1, \dots, Y_k]$$

denote the (unique) map of A -algebras with

$$\varphi(eT) = f(e) + \sum_{i=1}^k f_i(e)Y_i$$

for $e \in E$. Note that φ is not homogeneous in general when we equip $\mathbf{S}(E)$ and $A[\underline{Y}]$ with their standard grading. In the following let $J := IA[\underline{Y}]$ be the extension ideal, $\mathfrak{M}$ the maximal homogeneous ideal $\mathfrak{m}A[\underline{Y}] + (\underline{Y})$ of $A[\underline{Y}]$ and $B := A[\underline{Y}]_{\mathfrak{M}}$ the localization. Obviously φ_f is the composed map

$$S \xrightarrow{\varphi} A[\underline{Y}] \xrightarrow{Y_i=0} A.$$

The next lemma is a standard fact; for completeness we give a short argument.

Lemma 3.5. *If for some prime $\mathfrak{p}$ of A the localized module $E_{\mathfrak{p}}$ is free of rank r , then the maps*

$$A_{\mathfrak{p}} \xrightarrow{\text{can}} S_{\mathfrak{p}} \xrightarrow{\varphi} A_{\mathfrak{p}}[\underline{Y}]$$

are polynomial ring extensions of relative dimension r and $k - r$, respectively.

Proof. Localizing at $\mathfrak{p}$ we may suppose that E is free over A and so $E^\vee$ has a basis, say $f_1, \dots, f_r$, where $r = \text{rk } E$. We can write $f = \sum_{\varrho=1}^r a_\varrho f_\varrho$ and $f_i = \sum_{\varrho=1}^r a_{i\varrho} f_\varrho$ for $i > r$ with $a_\varrho, a_{i\varrho} \in A$. The basis $f_1, \dots, f_r$ provides an isomorphism between the symmetric algebra S and the polynomial ring $A[Y_1, \dots, Y_r]$, and the map $S \cong A[Y_1, \dots, Y_r] \rightarrow A[\underline{Y}]$ is given by $Y_\varrho \mapsto a_\varrho + Y_\varrho + \sum_{i>r} a_{i\varrho} Y_i$, $1 \leq \varrho \leq r$. Hence via this map, $A[\underline{Y}]$ may be considered as a polynomial ring over S in the indeterminates $Y_{r+1}, \dots, Y_k$. $\square$

The following lemma is a crucial step in the proof of Theorems 3.3 and 3.10; it is the only place where we use the assumption (**).

Lemma 3.6. *With the notation of 3.4 and the assumptions of Theorem 3.3 the following hold :*

- (1) *All minimal primes of $J \subseteq A[\underline{Y}]$ are contained in $\mathfrak{M}$.*
- (2) *Assume moreover that $A = \hat{A}$ is complete and E has an isolated singularity. If $S/I : \mathfrak{m}^\infty$ is an integral domain then so is the completion of $B/JB : \mathfrak{m}^\infty$.*

Proof. As $f \in \mathfrak{m}E^\vee$ we can write $f = \sum_i t_i f_i$ with $t_i \in \mathfrak{m}$. Let $\psi : A[\underline{Y}] \rightarrow A[\underline{Y}]$ be the isomorphism of A -algebras given by $Y_i \mapsto Y_i - t_i$. The composition

$$\psi \circ \varphi : S \longrightarrow A[\underline{Y}]$$

maps eT onto $f(e) + \sum f_i(e)(Y_i - t_i) = \sum f_i(e)Y_i$ and so $\psi \circ \varphi$ is homogeneous. The homogeneous ideal $\psi \circ \varphi(I)A[\underline{Y}] = \psi(J)$ has only homogeneous minimal primes that are all contained in $\mathfrak{M}$. As ψ maps $\mathfrak{M}$ into itself, (1) follows.

In order to prove (2) we first notice that for every prime $\mathfrak{p} \neq \mathfrak{m}$ of A the map $\psi \circ \varphi : S_{\mathfrak{p}}/I_{\mathfrak{p}} \rightarrow A_{\mathfrak{p}}[\underline{Y}]/\psi(J_{\mathfrak{p}})$ is a polynomial ring extension by Lemma 3.5. Thus it follows that $A[\underline{Y}]/(\psi(J) : \mathfrak{m}^\infty)$ is a homogeneous domain and so its completion, which is equal to the completion of $B/(\psi(J)B : \mathfrak{m}^\infty)$, is a domain. As the rings $B/(JB : \mathfrak{m}^\infty)$ and $B/(\psi(J)B : \mathfrak{m}^\infty)$ are isomorphic under the map induced by ψ , (2) follows. $\square$

Proof of Theorem 3.3. We may assume that $A = \hat{A}$ is complete. Furthermore, if $\dim(A/(\mathfrak{q} + I(f))) \geq \dim A/\mathfrak{q} - \tau$, then $\dim A/I(f) \geq \dim A - \tau$. Thus, replacing A by $A/\mathfrak{q}$ we are reduced to the case where A is a complete domain and $\mathfrak{q} = 0$. Let $\mathfrak{p}$ be a minimal prime of I that extends to a minimal prime of $I_{\mathfrak{q}}$ of maximal dimension in $S_{\mathfrak{q}}$. Clearly it suffices to show the result for $\mathfrak{p}$ instead of I . Thus we may further suppose that I is prime.

According to Lemma 3.5, $S_{\mathfrak{q}}/I_{\mathfrak{q}} \rightarrow A_{\mathfrak{q}}[\underline{Y}]/J_{\mathfrak{q}}$ is a polynomial extension of relative dimension $k - r$. Thus there is a minimal prime $\mathfrak{P}$ of J contracting to I . Lemma 3.6(1) shows that $\mathfrak{P}$ is contained in $\mathfrak{M}$. Applying the dimension formula [EGA, 5.5.8.1] for morphisms of finite type we get

$$\dim B/\mathfrak{P}B = \dim A + \text{trdeg}_{A_{\mathfrak{q}}} B_{\mathfrak{q}}/\mathfrak{P}B_{\mathfrak{q}}.$$

By Lemma 3.5, $\text{trdeg}_{A_{\mathfrak{q}}} B_{\mathfrak{q}}/\mathfrak{P}B_{\mathfrak{q}} = \text{trdeg}_{A_{\mathfrak{q}}} S_{\mathfrak{q}}/I_{\mathfrak{q}} + (k - r) = k - \tau$, where we used our assumption that $\dim S_{\mathfrak{q}}/I_{\mathfrak{q}} = r - \tau$. Thus we obtain

$$\dim B/JB \geq \dim B/\mathfrak{P}B = \dim A + k - \tau.$$

Using Krull's principal ideal theorem it follows that

$$B/(JB + (Y_1, \dots, Y_k)) \cong A/I(f)$$

has dimension $\geq \dim A - \tau$, as desired. $\square$

Example 3.7. Let $(A, \mathfrak{m})$ be the localization of the ring $R := \mathbb{C}[X_0, \dots, X_3]$ at the homogeneous maximal ideal. Consider the A -module $E := \text{coker}(h : A \rightarrow \Theta_A)$, where Θ_A is the module of derivations on A and h is given by $h(1) = \sum X_i \partial / \partial X_i$. The element $f := X_0 dX_1 - X_1 dX_0 + X_2 dX_3 - X_3 dX_2 \in E^\vee$ has then the property $f(E) = \mathfrak{m}$. Taking $I \subseteq \mathbf{S}(E)$ to be the ideal generated by the forms of strictly positive degree we have $\dim \mathbf{S}(E_{\mathfrak{q}})/I_{\mathfrak{q}} = 0 = \text{rk } E - 3$ with $\mathfrak{q} := 0$ the minimal prime of A , but $\dim A/I(f) = 0 < \dim A - 3$. This example shows that the assumption $f \in \mathfrak{m}E^\vee$ is essential.

It is instructive to understand geometrically why Lemma 3.6(1) does not hold in this example. The ring A can be regarded as the local ring of the affine 4-space $V := \text{Spec } R$ at the origin $0 \in V$ and E is the localization of $E_R := \text{coker}(h : R \rightarrow \Theta_R)$ with h as above. Thus $\mathbb{V}(E_R) := \text{Spec } \mathbf{S}(E_R)$ embeds into the affine space $V^\vee \times V = \mathbb{V}(\Theta_R)$. Taking as generators of $E_R^\vee$ the standard syzygies $f_{ij} = X_i dX_j - X_j dX_i$ we have $\text{Spec } R[\underline{Y}] \cong \Lambda^2 V^\vee \times V := \mathbb{V}(\Lambda^2 \Theta_R)$, and the generators f_{ij} induce a morphism $H : \Lambda^2 V^\vee \times V \rightarrow \mathbb{V}(E_R)$. The composition with the embedding $\mathbb{V}(E_R)$ into $V^\vee \times V$ on the $\mathbb{C}$ -valued points is the map

$$\begin{aligned} \Lambda^2 V^\vee \times V &\longrightarrow \mathbb{V}(E_R) \longrightarrow V^\vee \times V \\ (T, x) &\mapsto (Tx, x). \end{aligned}$$

Here we consider a $\mathbb{C}$ -valued point of $\Lambda^2 V^\vee$ as a skew symmetric matrix T . The given element f corresponds to a nondegenerate matrix T_0 and the map φ in 3.4 to $\Phi : (T, x) \mapsto ((T + T_0)x, x)$. The ideal I defines the zero section $\{0\} \times V$, and the preimage of this zero section in $\Lambda^2 V^\vee \times V$ is the set

$$Z := \{(T, x) : (T + T_0)x = 0\}.$$

The irreducible components of Z are given by

$$Z_1 := \{(T, x) : (T + T_0)x = 0 \text{ and } \text{Pf}(T + T_0) = 0\} \quad \text{and} \quad Z_2 := V^\vee \times \{0\},$$

where Z_1 is of codimension 3 and Z_2 of codimension 4. Thus Z_1 does not meet $\{0\} \times V$, and the hypersurfaces $Y_{ij} = 0$ intersect Z in the origin $(0, 0)$.

Remark 3.8. In the setting of Theorem 3.2 there is another condition (***) that implies (*), but is weaker than assumption (**) in Theorem 3.3. We abbreviate as before $-\times = \text{Hom}_{A'}(-, A')$. If

$$(***) \quad f \otimes 1 \in \mathfrak{m}'(E^\vee \otimes_A A')^{\times \times},$$

then (*) holds. This implication follows from the existence of the natural A' -linear map

$$(E^\vee \otimes_A A')^{\times \times} \longrightarrow (E \otimes_A A')^{\times \times \times} \longrightarrow (E \otimes_A A')^\times \cong \text{Hom}_A(E, A'),$$

that sends the image of $f \otimes 1$ to $h \circ f$.

Condition (***) for h any high power of the Frobenius map is precisely the p -ampleness of the pair $(Af, E^\vee)$ as introduced in Definition 2.2. Thus we deduce:

Theorem 3.9. *Assume that $(A, \mathfrak{m})$ is a noetherian local ring of prime characteristic p . The conclusion of Theorem 3.3 remain valid if one replaces assumption (**) by the weaker condition that the pair*

$$(***) \quad (Af, E^\vee) \quad \text{is } p\text{-ample.}$$

We now turn to our results about connectedness.

Theorem 3.10. *Let $(A, \mathfrak{m})$ be a noetherian local ring, E a finite A -module of rank r with an isolated singularity, and $f \in E^\vee$. Assume that*

$$(*) \quad h \circ f \in \mathfrak{m}' \operatorname{Hom}_A(E, A')$$

for some local homomorphism $h : A \rightarrow A'$ with $(A', \mathfrak{m}')$ a noetherian local ring so that the map of completions $\hat{h} : \hat{A} \rightarrow \hat{A}'$ has nilpotent kernel and $\hat{A}'$ is integral over the image of $\hat{h}$. Let $I \subseteq S := \mathbf{S}(E)$ be a homogeneous ideal and suppose that the map

$$\operatorname{Spec}(S/I) \setminus V(\mathfrak{m}(S/I)) \longrightarrow \operatorname{Spec}(A) \setminus \{\mathfrak{m}\}$$

has non empty geometrically irreducible fibers all of the same dimension $r - \tau$. If $\hat{A}$ is connected in dimension d then $\hat{A}/I(f)\hat{A}$ is connected in dimension $d - \tau$, where $I(f) := \varphi_f(I)$.

Once again, the special case $h = \operatorname{id}_A$ is noteworthy:

Theorem 3.11. *Let $(A, \mathfrak{m})$ be a noetherian local ring, E a finite A -module of rank r with an isolated singularity, and assume that*

$$(**) \quad f \in \mathfrak{m}E^\vee.$$

Let $I \subseteq S := \mathbf{S}(E)$ be a homogeneous ideal and suppose that the map

$$\operatorname{Spec}(S/I) \setminus V(\mathfrak{m}(S/I)) \longrightarrow \operatorname{Spec}(A) \setminus \{\mathfrak{m}\}$$

has non empty geometrically irreducible fibers all of the same dimension $r - \tau$. If $\hat{A}$ is connected in dimension d then $\hat{A}/I(f)\hat{A}$ is connected in dimension $d - \tau$, where $I(f) := \varphi_f(I)$.

In the light of Remark 3.8, applying Theorem 3.10 to the iterated Frobenius $h = F^a$, $a \gg 0$, gives:

Theorem 3.12. *Assume that $(A, \mathfrak{m})$ is a noetherian local ring of prime characteristic p . The conclusion of Theorem 3.10 remains valid if one replaces assumption $(**)$ by the weaker condition that the pair*

$$(***) \quad (Af, E^\vee) \text{ is } p\text{-ample}.$$

To prove Theorem 3.10 we will need the next two lemmas.

Lemma 3.13. *Let Y be a noetherian scheme and let $Y_i \subseteq Y$ be closed subsets of dimension $\geq d + 1$ with $Y = Y_1 \cup \dots \cup Y_s$. Consider the following conditions :*

- (i) *Y is connected in dimension d .*
- (ii) *For all Y_i, Y_j there exists a sequence $Y_{i_1}, \dots, Y_{i_t}$ such that $Y_i = Y_{i_1}$, $Y_j = Y_{i_t}$ and*

$$\dim(Y_{i_\alpha} \cap Y_{i_{\alpha+1}}) \geq d \text{ for } 1 \leq \alpha < t.$$

Then (i) implies (ii). If all Y_i are connected in dimension d then also the converse is true.

Proof. For (ii) $\Rightarrow$ (i) we refer the reader to [FOV, Proposition 3.1.4]. To prove (i) $\Rightarrow$ (ii), note first that the relation given by $i \sim j$ if and only if (ii) is satisfied, is an equivalence relation. For an equivalence class $\alpha \subseteq \{1, \dots, s\}$ let $Y_\alpha := \bigcup_{i \in \alpha} Y_i$. Clearly $Y = \bigcup_\alpha Y_\alpha$ and $\dim(Y_\alpha \cap Y_\beta) < d$ for $\alpha \neq \beta$. Thus Y will be not connected in dimension d unless there is only one equivalence class. $\square$

Lemma 3.14. *Let Y be an irreducible noetherian scheme and $\mathcal{R}$ a finitely generated positively graded $\mathcal{O}_Y$ -algebra with coherent homogeneous components. If the morphism $X := \operatorname{Spec} \mathcal{R} \rightarrow Y$ has irreducible fibers all of the same dimension, then X is irreducible.*

Proof. If $\mathcal{R}$ is a finite $\mathcal{O}_Y$ -module then the assertion is immediate. Otherwise the morphism

$$\pi : Z := \operatorname{Proj} \mathcal{R} \rightarrow Y$$

has non empty irreducible fibers all of the same dimension, say $s \geq 0$. Let Z_0 be an irreducible component of Z with generic point η_0 such that $\pi(Z_0) = Y$. The generic fiber $F := \pi^{-1}(\pi(\eta_0))$ through η_0 contains $F \cap Z_0$ as an irreducible component and so these sets are equal by our assumption. By the dimension formula $\dim Z_0 = s + \dim Y$. Using the semicontinuity of fiber dimensions all fibers of $Z_0 \rightarrow Y$ have dimension $\geq s$. Therefore they are equal to the fibers of π as these are irreducible of dimension s by our assumption. Hence $Z = Z_0$ and so Z and then also X are irreducible, as desired. $\square$

Proof of Theorem 3.10. We may assume that $A = \hat{A}$ and $A' = \hat{A}'$ are complete. We may further suppose that $d > 0$. Let Y_i be the irreducible components of $Y = \operatorname{Spec} A$ and suppose that we can prove that $V_i := Y_i \cap V$ is connected in dimension $d - \tau$, where $V := V(I(f))$. We claim that then V is connected in dimension $d - \tau$. Indeed, applying Lemma 3.13, for all Y_i, Y_j there exists a sequence of irreducible components $Y_i = Y_{i_1}, \dots, Y_{i_t} = Y_j$, with $\dim(Y_{i_\alpha} \cap Y_{i_{\alpha+1}}) \geq d$ for $1 \leq \alpha < t$. Taking the intersection with V and using Theorem 3.2 we obtain that $\dim(V_{i_\alpha} \cap V_{i_{\alpha+1}}) \geq d - \tau$ for $1 \leq \alpha < t$. Hence again by Lemma 3.13, V is connected in dimension $d - \tau$.

As $h : A \rightarrow A'$ has nilpotent kernel and A' is integral over the image of h , the induced map $\operatorname{Spec} A' \rightarrow \operatorname{Spec} A$ is surjective. Thus, by factoring out an arbitrary minimal prime of A and a prime of A' lying over it, we may suppose that $A \subseteq A'$ is an integral extension of domains. As $\dim A' = \dim A > d$ it follows that A' is connected in dimension d . With $I'(f')$ defined as in the proof of Theorem 3.2 one has $I'(f') = I(f)A'$. Since $A \subseteq A'$ is an integral extension, connectedness in dimension d descends from $A'/I(f)A'$ to $A/I(f)$. Replacing A by A' we may then assume that $A = A'$ and that the stronger condition $(**)$ of Theorem 3.11 holds. Furthermore A is a complete domain.

After replacing the ideal I by its radical, Lemma 3.14 shows that $S/I : \mathfrak{m}^\infty$ is a domain, and then by Lemma 3.6(2) the same is true for the completion of $B/JB : \mathfrak{m}^\infty$. Applying Grothendieck's connectedness theorem we obtain that

$$V(JB : \mathfrak{m}^\infty + (Y_1, \dots, Y_k)) \subseteq \operatorname{Spec} B/(Y_1, \dots, Y_k) \cong \operatorname{Spec} A$$

is connected in dimension $d - \tau$. As this set is equal to $V(I(f))$ the result follows. $\square$

We can also generalize our results in the spirit of [EHU₃, Theorem 3.1]. For instance:

Corollary 3.15. *Theorem 3.3 remains valid if we replace condition $(**)$ by the weaker condition that*

*$(**')$ there exists a local map $h : A \rightarrow A'$ to a noetherian local ring $(A', \mathfrak{m}')$ so that $\mathfrak{q} = h^{-1}(\mathfrak{q}')$ for some prime $\mathfrak{q}'$ of A' and $f \otimes 1 \in \mathfrak{m}'(E^\vee \otimes_A A'/\text{torsion})$.*

Proof. Inspecting the proof of Theorem 3.3 we only need to show the following analogue of Lemma 3.6(1): All minimal primes $\mathfrak{P}$ of $J \subseteq A[\underline{Y}]$ with $\mathfrak{P} \cap A = \mathfrak{q}$ are contained in $\mathfrak{M}$.

As $f \otimes 1 \in \mathfrak{m}'(E^\vee \otimes_A A'/\text{torsion})$ we can write $f \otimes 1 = \sum_i t_i f_i \otimes 1$ with $t_i \in \mathfrak{m}'$. Let $\psi : A[\underline{Y}] \rightarrow A'[\underline{Y}]$ be the morphism given by $Y_i \mapsto Y_i - t_i$. As in the proof of Lemma 3.6(1) the composition $\psi \circ \varphi : S \rightarrow A'[\underline{Y}]$ is homogeneous. The homogeneous ideal $\psi \circ \varphi(I)A'[\underline{Y}] = \psi(J)A'[\underline{Y}]$ has only homogeneous minimal primes that are contained in the maximal homogeneous ideal $\mathfrak{M}'$ of $A'[\underline{Y}]$. Moreover ψ maps $\mathfrak{M}$ into $\mathfrak{M}'$. Since $A_{\mathfrak{q}}$ is Artinian and the map $A_{\mathfrak{q}} \rightarrow A'_{\mathfrak{q}'}$ is local, it follows that every minimal prime of $JA_{\mathfrak{q}}[\underline{Y}] \subseteq A_{\mathfrak{q}}[\underline{Y}]$ is contracted from a minimal prime of $\psi(J)A'_{\mathfrak{q}'}[\underline{Y}] \subseteq A'_{\mathfrak{q}'}[\underline{Y}]$. Thus for every minimal prime $\mathfrak{P}$ of $J \subseteq A[\underline{Y}]$ with $\mathfrak{P} \cap A = \mathfrak{q}$ there exists a minimal prime $\mathfrak{P}'$ of $\psi(J)A'[\underline{Y}] \subseteq A'[\underline{Y}]$ lying over it. As $\mathfrak{P}' \subseteq \mathfrak{M}'$ we conclude that $\mathfrak{P} \subseteq \mathfrak{M}$. $\square$

4. CODIMENSION AND CONNECTEDNESS OF DEGENERACY LOCI

4.1. Maps of modules. While it is clear how to define determinantal ideals and degeneracy loci for morphisms of locally free sheaves, for maps of arbitrary coherent sheaves several definitions are possible. For our purposes the following natural notion is best suited.

4.1. Given a morphism $f : \mathcal{M} \rightarrow \mathcal{N}$ of coherent sheaves on a scheme X we can introduce the $(t+1)$ -st *determinantal ideal* of f by

$$\mathcal{I}_{t+1}(f) := \text{Im}(\Lambda^{t+1} \mathcal{M} \otimes_{\mathcal{O}_X} \Lambda^{t+1} \mathcal{N}^\vee \longrightarrow \mathcal{O}_X),$$

where the map is defined by

$$m_0 \otimes \dots \otimes m_t \otimes \varphi_0 \otimes \dots \otimes \varphi_t \longmapsto \det(\varphi_i \circ f(m_j))$$

for local sections $m_0, \dots, m_t$ in $\mathcal{M}$ and $\varphi_0, \dots, \varphi_t$ in $\mathcal{N}^\vee$. Its set of zeros $D_t(f) := V(\mathcal{I}_{t+1}(f))$ will be called the t -th *degeneracy locus* of f .

Similarly, if $f : M \rightarrow N$ is a linear map of A -modules then $\mathcal{I}_{t+1}(f)$ denotes the image of the induced map $\Lambda^{t+1} M \otimes_A \Lambda^{t+1} N^\vee \longrightarrow A$ and $D_t(f) := V(\mathcal{I}_{t+1}(f)) \subseteq \text{Spec } A$ the t -th degeneracy locus. Note that the sheaf $\tilde{\mathcal{I}}_{t+1}(f)$ associated to $\mathcal{I}_{t+1}(f)$ on $X = \text{Spec } A$ is just the $(t+1)$ -st determinantal ideal associated to $\tilde{f} : \tilde{M} \rightarrow \tilde{N}$.

If $\mathcal{M}$ and $\mathcal{N}$ are free $\mathcal{O}_X$ -modules of rank m and n , respectively, then f can be described by a $n \times m$ -matrix F . In this case $\mathcal{I}_{t+1}(f) = \mathcal{I}_{t+1}(F)$ is just the ideal generated by the $(t+1)$ -minors of F .

Theorem 4.2. *Let $(A, \mathfrak{m})$ be a noetherian local ring and $f : M \rightarrow N$ a linear map of finite A -modules. Assume there exists a prime ideal $\mathfrak{q} \subseteq A$ with $\dim A = \dim A/\mathfrak{q}$ such that $M_{\mathfrak{q}}$ and $N_{\mathfrak{q}}$ are free $A_{\mathfrak{q}}$ -modules of ranks m and n , respectively. If $t \leq \min\{n, m\}$ then with $\tau := (n-t)(m-t)$ we have $\dim D_t(f) \geq \dim A - \tau$ in each of the following cases :*

- (a) $f \in \mathfrak{m} \text{Hom}(M, N)$.
- (b) A has prime characteristic p and the pair $(Af, \text{Hom}(M, N))$ is p -ample.

Proof. With $E := \text{Hom}(M, N)^\vee$ the morphism f can be considered as an element in $E^\vee$. The idea is to apply Theorems 3.3 and 3.9 to the “generic” determinantal ideal $\mathcal{I}_{t+1} \subseteq S := \mathbf{S}(E)$.

In the case that N is free the definition of I_{t+1} is easy: the evaluation map $\text{Hom}(M, N) \otimes_A M \rightarrow N$ induces a map $M \rightarrow \text{Hom}(M, N)^\vee \otimes_A N = E \otimes_A N$. It gives a “universal” morphism

$$F : \mathbf{S}(E) \otimes_A M \longrightarrow \mathbf{S}(E) \otimes_A N,$$

and one can take $I_{t+1} := I_{t+1}(F)$. If N is not free no such universal map exists, but one can nevertheless define the generic determinantal ideal as follows:

Applying the evaluation map twice yields a map $M \otimes_A \text{Hom}(M, N) \otimes_A N^\vee \rightarrow A$, which induces

$$\epsilon : M \otimes_A N^\vee \longrightarrow E = \text{Hom}(M, N)^\vee \subseteq S = \mathbf{S}(E).$$

We can now define the “generic” determinantal ideal $I_{t+1} \subseteq S$ to be the ideal generated by the image of the map

$$\begin{aligned} \Lambda^{t+1} M \otimes_A \Lambda^{t+1}(N^\vee) &\longrightarrow S \\ m_0 \wedge \dots \wedge m_t \otimes \varphi_0 \wedge \dots \wedge \varphi_t &\longmapsto \det(\epsilon(m_j \otimes \varphi_i)) \in S. \end{aligned}$$

It is clear that

- (A) the construction yields the previously defined determinantal ideal (see 4.1) when N is free; in particular I_{t+1} is the ideal of $(t+1)$ -minors of an $n \times m$ matrix of variables if both M and N are free of ranks m and n , respectively;
- (B) applying the map $\varphi_f : S \rightarrow A$ induced by f (see (3.1), one obtains $\varphi_f(I_{t+1}) = I_{t+1}(f)$, the determinantal ideal of f .

Also notice that $I_{t+1} \subseteq S$ is a homogeneous ideal generated by forms of positive degree. From item (A) and Lemma 4.3 below we see that $\dim S_{\mathbf{q}}/I_{\mathbf{q}} = r - \tau$ with $r := nm = \text{rank } E_{\mathbf{q}}$. As $I_{t+1}(f) = \varphi_f(I_{t+1})$ by item (B), the present theorem now follows from Theorem 3.3 in case (a) and Theorem 3.9 in case (b). $\square$

Lemma 4.3. *Let $S = A[T_{ij}]$ be the polynomial ring over a noetherian ring in the variables T_{ij} , $1 \leq i \leq n$, $1 \leq j \leq m$, and let I_{t+1} be the ideal of $(t+1)$ -minors of the generic matrix $T = (T_{ij})$, where $t \leq \min\{n, m\}$. Then S/I_{t+1} is flat over A with geometrically normal fibers of dimension $nm - \tau$, where $\tau := (n-t)(m-t)$.*

Likewise this holds for the generic symmetric $m \times m$ -matrix, in which case the fiber dimension is

$$\binom{m+1}{2} - \tau \quad \text{with } \tau := \binom{m-t+1}{2},$$

and for the generic alternating $m \times m$ -matrix, for which the fiber dimension is

$$\binom{m}{2} - \tau \quad \text{with } t := 2s \text{ and } \tau := \binom{m-2s}{2}.$$

For the proofs we refer the reader to [EN] in the generic case, [Jo, 2.1 and 2.4] in the symmetric case and [JP, 2.1 and 2.5] in the alternating case.

Theorem 4.4. *Let $(A, \mathfrak{m})$ be a noetherian local ring and let $f : M \rightarrow N$ be a linear map of finite A -modules with an isolated singularity of ranks m and n , respectively. Assume that one of the conditions (a), (b) in Theorem 4.2 is satisfied. If $\hat{A}$ is connected in dimension d and $t \leq \min\{n, m\}$, then with $\tau := (n-t)(m-t)$ the degeneracy locus $D_t(f)$ is connected in dimension $d - \tau$.*

Proof. As in the proof of Theorem 4.2 we let E be the dual of $\text{Hom}(M, N)$ and $I = I_{t+1} \subseteq \mathbf{S}(E)$ the “generic” determinantal ideal so that $\varphi_f(I) = I_{t+1}(f)$ in the notation of Theorems 3.11 and 3.12. The map $V(I) \rightarrow \text{Spec } A$ is flat over points $\mathfrak{p} \neq \mathfrak{m}$, and the fibers are generic determinantal varieties. According to Lemma 4.3 these fibers are geometrically normal of constant dimension $nm - \tau$. Now the result follows from Theorems 3.11 and 3.12. $\square$

Order ideals are a special case of determinantal ideals, namely the case where $M = A$ and $t = 0$. Applying Theorems 4.2 and 4.4 to this case we obtain for instance the following result:

Corollary 4.5. *Let $(A, \mathfrak{m})$ be a noetherian local ring and let N be a finite A -module with an isolated singularity of rank n . Assume that $f \in N$ satisfies one the the following two conditions :*

- (a) $f \in \mathfrak{m}N$.
- (b) A has prime characteristic $p > 0$ and the pair (Af, N) is p -ample.

Then the following hold :

- (1) $\dim(A/N^\vee(f)) \geq \dim A - n$.
- (2) *If $\hat{A}$ is connected in dimension d then $A/N^\vee(f)$ is connected in dimension $d - n$.*

Remark 4.6. (1) One should expect that Theorems 4.2 and 4.4 remain valid in arbitrary characteristic if the pair $(Af, \text{Hom}(M, N))$ is assumed to be ample. A positive answer to this problem would yield the corresponding projective results (see Remark 4.16(1)) in full generality.

(2) We do not know whether in Theorems 4.2 and 4.4 one can also obtain the refined bounds of Bruns [Br] for determinantal ideals $I_{t+1}(f)$ when $I_{t+2}(f) = 0$. We note that the analogous question for determinantal ideal of symmetric maps is not true in positive characteristic as is seen by the simple example of the symmetric matrix

$$F = \begin{pmatrix} 0 & X_1 & X_2 \\ X_1 & 0 & X_3 \\ X_2 & X_3 & 0 \end{pmatrix}$$

in characteristic 2, see [EHU₄]. Therefore one cannot expect that the solution to this problem follows from the results of Section 3.

4.2. Symmetric maps. Let M and L be finite modules over the noetherian local ring $(A, \mathfrak{m})$. In this subsection we will study degeneracy loci and determinantal ideals of symmetric homomorphisms $f : M \rightarrow \text{Hom}_A(M, L)$. By this we mean that the corresponding map in $\text{Hom}_A(M \otimes_A M, L)$ factors through the symmetric power $\mathbf{S}^2(M)$. By $\text{Hom}^s(M, \text{Hom}(M, L))$ we denote the A -module of all such symmetric maps so that

$$\text{Hom}^s(M, \text{Hom}(M, L)) \cong \text{Hom}(\mathbf{S}^2 M, L)$$

With this notation we have the following result:

Theorem 4.7. *Assume there exists a prime ideal $\mathfrak{q} \subseteq A$ with $\dim A = \dim A/\mathfrak{q}$ such that $M_{\mathfrak{q}}$ and $L_{\mathfrak{q}}$ are free of ranks m and 1, respectively. If $t \leq m$ then with $\tau := \binom{m-t+1}{2}$ we have $\dim D_t(f) \geq \dim A - \tau$ in each of the following cases :*

- (a) $f \in \mathfrak{m} \text{Hom}^s(M, \text{Hom}(M, L))$.
- (b) A has prime characteristic p and the pair $(Af, \text{Hom}^s(M, \text{Hom}(M, L)))$ is p -ample.

Proof. With $E := \operatorname{Hom}^s(M, \operatorname{Hom}(M, L))^\vee$ the morphism f can be considered as an element in $E^\vee$. As in the proof of Theorem 4.2 the idea is to apply Theorems 3.3 and 3.9 to the “generic” symmetric determinantal ideal $I_{t+1} \subseteq S := \mathbf{S}(E)$, which can be defined as follows:

Applying the evaluation map twice yields a map

$$M \otimes_A \operatorname{Hom}^s(M, \operatorname{Hom}(M, L)) \otimes_A \operatorname{Hom}(M, L)^\vee \longrightarrow A,$$

which in turn induces

$$\epsilon : M \otimes_A \operatorname{Hom}(M, L)^\vee \longrightarrow E = \operatorname{Hom}^s(M, \operatorname{Hom}(M, L))^\vee \subseteq S = \mathbf{S}(E).$$

We then define the “generic” symmetric determinantal ideal $I_{t+1} \subseteq S$ to be the ideal generated by the image of the map

$$\begin{aligned} \Lambda^{t+1} M \otimes_A \Lambda^{t+1}(\operatorname{Hom}(M, L)^\vee) &\longrightarrow S \\ m_0 \wedge \dots \wedge m_t \otimes \varphi_0 \wedge \dots \wedge \varphi_t &\longmapsto \det(\epsilon(m_j \otimes \varphi_i)) \in S. \end{aligned}$$

One easily sees that

- (A) I_{t+1} is the ideal of $(t+1)$ -minors of a symmetric $m \times m$ matrix of variables if both M and L are free of ranks m and 1 , respectively;
- (B) applying the map $\varphi_f : S \rightarrow A$ induced by f (see 3.1), one obtains the ideal $\varphi_f(I_{t+1}) = I_{t+1}(f)$ defining the degeneracy locus $D_t(f)$.

Furthermore $I_{t+1} \subseteq S$ is a homogeneous ideal generated by forms of positive degree. From property (A) and Lemma 4.3 one sees that $\dim S_{\mathfrak{q}}/I_{\mathfrak{q}} = r - \tau$ with $r := \binom{m+1}{2} = \operatorname{rank} E_{\mathfrak{q}}$. Since $I_{t+1}(f) = \varphi_f(I_{t+1})$ by property (B), the theorem follows once again from Theorem 3.3 in case (a) and Theorem 3.9 in case (b). $\square$

Theorem 4.8. *Assume in Theorem 4.7 that M and L are modules with an isolated singularity of ranks m and 1 , respectively, and that one of the conditions (a) or (b) is satisfied. If $\hat{A}$ is connected in dimension d and $t \leq m$, then with $\tau := \binom{m-t+1}{2}$ the degeneracy locus $D_t(f)$ is connected in dimension $d - \tau$.*

Proof. The proof is analogous to that of Theorem 4.4. $\square$

4.3. Alternating maps. Let M and L be finite modules over the noetherian local ring $(A, \mathfrak{m})$. The results of the preceding section hold likewise for alternating maps $f : M \rightarrow \operatorname{Hom}_A(M, L)$, i.e. maps corresponding to elements in $\operatorname{Hom}_A(\Lambda^2 M, L)$. By $\operatorname{Hom}^a(M, \operatorname{Hom}(M, L))$ we denote the A -module of all such alternating maps. The proofs of the next two results are analogous to those in the symmetric case and are left to the reader.

Theorem 4.9. *Assume there exists a prime ideal $\mathfrak{q} \subseteq A$ with $\dim A = \dim A/\mathfrak{q}$ such that $M_{\mathfrak{q}}$ and $L_{\mathfrak{q}}$ are free $A_{\mathfrak{q}}$ -modules of ranks m and 1 respectively. With $t := 2s \leq m$ and $\tau := \binom{m-2s}{2}$ we have $\dim D_t(f) \geq \dim A - \tau$ in each of the following cases $\therefore$.*

- (a) $f \in \mathfrak{m} \operatorname{Hom}^a(M, \operatorname{Hom}(M, L))$.
- (b) A has prime characteristic p and the pair $(Af, \operatorname{Hom}^a(M, \operatorname{Hom}(M, L)))$ is p -ample.

Theorem 4.10. *Assume in Theorem 4.9 that M and L are modules with an isolated singularity of ranks m and 1 , respectively, and that one of the conditions (a) or (b) is satisfied. If $\hat{A}$ is connected in dimension d and $t := 2s \leq m$, then with $\tau := \binom{m-2s}{2}$ the degeneracy locus $D_t(f)$ is connected in dimension $d - \tau$.*

4.4. Further results. One can generalize Theorems 3.10, 3.11, 3.12, 4.4, 4.8 and 4.10 to the case where the modules do not have an isolated singularity. As a sample we state the following result:

Corollary 4.11. *Let $(A, \mathfrak{m})$ be a noetherian local ring, $\mathfrak{a} \subseteq A$ an ideal, and $f : M \rightarrow N$ be a map of finite A -modules that are locally free on $\text{Spec } A \setminus V(\mathfrak{a})$ of constant ranks m and n , respectively. Assume that with $t \leq \min\{m, n\}$ and $\tau := (m-t)(n-t)$ we have $\dim A/\mathfrak{a} < \dim A - \tau$. If one of the conditions (a) or (b) in Theorem 4.2 is satisfied and $\hat{A}$ is connected in dimension d , then $D_t(f) \setminus V(\mathfrak{a})$ is connected in dimension $d - 1 - \tau$.*

Proof. We may assume that A is complete, and proceed by induction on $\dim A/\mathfrak{a}$. In case $\dim A/\mathfrak{a} = 0$ this is just Theorem 4.4. Now let $\dim A/\mathfrak{a} > 0$. By Theorem 4.2 $D_t(f) \setminus V(\mathfrak{a})$ is not empty and hence has dimension $> d - 1 - \tau$. Thus if $D_t(f) \setminus V(\mathfrak{a})$ is not connected in dimension $d - 1 - \tau$, then there exist proper closed subsets V_1, V_2 of $D_t(f) \setminus V(\mathfrak{a})$ with $D_t(f) \setminus V(\mathfrak{a}) = V_1 \cup V_2$ and $\dim V_1 \cap V_2 < d - 1 - \tau$. For a sufficiently general element $x \in \mathfrak{m}$ we have

$$\dim A/(xA + \mathfrak{a}) = \dim A/\mathfrak{a} - 1 < \dim A/xA - \tau$$

and $\dim V_1 \cap V_2 \cap V(x) < d - 2 - \tau$. Moreover, since every irreducible component of $D_t(f) \setminus V(\mathfrak{a})$ has dimension $\geq d - \tau$, the subsets $V_i \cap V(x)$ of $D_t(f) \cap V(x) \setminus V(\mathfrak{a})$ are proper. Hence $D_t(f) \cap V(x) \setminus V(\mathfrak{a})$ is not connected in dimension $d - 2 - \tau$. On the other hand, A/xA is connected in dimension $d - 1$ by Grothendieck's connectedness theorem, and conditions (a) and (b) pass from A to A/xA according to Remark 2.3(4). Therefore the induction hypothesis shows that $D_t(f) \cap V(x) \setminus V(\mathfrak{a})$ is connected in dimension $d - 2 - \tau$, which is a contradiction. $\square$

Remark 4.12. Similar results hold for symmetric and alternating maps. We leave the straightforward formulations and proofs to the reader.

One can also generalize the preceding results in the spirit of Theorems 3.3, 3.10 and Corollary 3.15. Again, we only give a sample of such a generalization. The special case of order ideals had been treated in [EHU₃].

Corollary 4.13. *Theorem 4.2 remains valid if we replace condition (a) by the weaker condition that there exists a local map $h : A \rightarrow A'$ to a noetherian local ring $(A', \mathfrak{m}')$ so that $\mathfrak{q} = h^{-1}(\mathfrak{q}')$ for some prime $\mathfrak{q}'$ of A' and*

$$f \otimes 1 \in \mathfrak{m}'(\text{Hom}_A(M, N) \otimes_A A'/\text{torsion}).$$

This is easily derived as in the proof of Theorem 4.2 using Corollary 3.15. Again we only mention the fact that similar generalizations hold for symmetric and alternating maps; the straightforward formulations and proofs are left to the reader.

Finally we apply these results to maps of vector bundles on projective schemes.

Corollary 4.14. *Let X be a projective K -scheme over field K of characteristic $p > 0$ and let $f : \mathcal{M} \rightarrow \mathcal{N}$ be an $\mathcal{O}_X$ -linear map of locally free sheaves of ranks m and n , respectively. If $\mathcal{H}om(\mathcal{M}, \mathcal{N})$ is cohomologically p -ample and $t \leq \min\{m, n\}$, then with $\tau := (m-t)(n-t)$ the following hold :*

- (1) *If $\dim X \geq \tau$ then $D_t(f) \neq \emptyset$.*
- (2) *If X is connected in dimension d then $D_t(f)$ is connected in dimension $d - \tau$.*

Proof. Write $X = \text{Proj } R$, where $R = K[R_1]$ is a finitely generated graded algebra over $K = R_0$. The R -modules

$$M = \bigoplus_{n \geq 0} H^0(X, \mathcal{M}(n)) \quad \text{and} \quad N = \bigoplus_{n \geq 0} H^0(X, \mathcal{N}(n))$$

are then finitely generated over R and $\mathcal{M} = \tilde{M}$, $\mathcal{N} = \tilde{N}$ are the sheaves associated to M , N , respectively. As $\mathcal{M}, \mathcal{N}$ are vector bundles on X , the modules M, N have an isolated singularity. Let $I_{t+1}(f_0) \subseteq R$ be the $(t+1)$ st determinantal ideal of the map $f_0 : M \rightarrow N$ induced by f . Clearly f_0 is an element of degree 0 in $\text{Hom}_R(M, N)$ and so by Proposition 2.4(1) and Remark 2.3(2) the pair $(Rf_0, \text{Hom}_R(M, N))$ is p -ample. Let $\mathfrak{m}$ be the homogeneous maximal ideal of R . Applying Theorem 4.2 to the local ring $A = R_{\mathfrak{m}}$ and the localized map $f_0 : M_{\mathfrak{m}} \rightarrow N_{\mathfrak{m}}$, we obtain that $\dim A/I_{t+1}(f_0)A \geq \dim A - \tau$ and so $\dim R/I_{t+1}(f_0)R \geq \dim R - \tau$. As $\mathcal{I}_{t+1}(f) \subseteq \mathcal{O}_X$ is the ideal sheaf induced by $I_{t+1}(f_0)$, the first part follows. Likewise part (2) is an easy consequence of Theorem 4.4. Here one also uses the fact that a projective scheme is connected in dimension d if and only if the local ring at the vertex of the affine cone is connected in dimension $d+1$ (see [FOV, Lemma 3.1.9]). $\square$

In the case of symmetric or alternating maps one can apply Theorems 4.7, 4.8, 4.9 and 4.10 instead. This leads to the next result. Here $\mathbf{D}^n$ denotes the n th divided power functor.

Corollary 4.15. *Let X is a projective K -scheme over a field K of characteristic $p > 0$. Assume that $\mathcal{M}$ is a locally free $\mathcal{O}_X$ -module of rank m and $\mathcal{L}$ is a line bundle on X . Then for $t \leq m$ the following hold :*

- (1) *If $f : \mathcal{M}^{\vee} \rightarrow \mathcal{M} \otimes_{\mathcal{O}_X} \mathcal{L}$ is a symmetric map and $\mathbf{D}^2 \mathcal{M} \otimes_{\mathcal{O}_X} \mathcal{L}$ is cohomologically p -ample, then statements (1) and (2) of Corollary 4.14 hold with $\tau := \binom{m-t+1}{2}$.*
- (2) *If $f : \mathcal{M}^{\vee} \rightarrow \mathcal{M} \otimes_{\mathcal{O}_X} \mathcal{L}$ is an alternating map and $\Lambda^2 \mathcal{M} \otimes_{\mathcal{O}_X} \mathcal{L}$ is cohomologically p -ample, then statements (1) and (2) of Corollary 4.14 hold with $t := 2s$ and $\tau := \binom{m-2s}{2}$.*

Remark 4.16. (1) Suppose that X is a projective variety over the field K . If K has characteristic 0 or X is smooth, the conclusion of Corollary 4.14 has been shown by Fulton and Lazarsfeld [FL₁] under the assumption that $\mathcal{H}om(\mathcal{M}, \mathcal{N})$ is ample. Similarly, Corollary 4.15 with the assumption of p -ampleness replaced by ampleness is due to Tu [Tu] provided that K has characteristic 0 and the maps have even ranks.

(2) Using Theorems 4.2 and 4.4 we also obtain an analogue of Corollary 4.14 in arbitrary characteristic if we replace the condition on the cohomological p -ampleness of the vector bundle $\mathcal{H} := \mathcal{H}om(\mathcal{M}, \mathcal{N})$ by the assumption that the element $f \in H^0(X, \mathcal{H})$ can be written as a linear combination of sections in $H^0(X, \mathcal{H}(-1))$ with coefficients in $H^0(X, \mathcal{O}_X(1))$. A similar remark applies to Corollary 4.15.

REFERENCES

- [Ba] Barton, C.: *Tensor products of ample bundles in characteristic p* . Amer. J. Math. **93**, 429–438 (1971)
- [Br] Bruns, W.: *The Eisenbud-Evans generalized principal ideal theorem and determinantal ideals*. Proc. Amer. Math. Soc. **83**, 19–24 (1981)
- [BV] Bruns, W.; Vetter, U.: *Determinantal rings*. Lecture Notes in Mathematics **1327**, Springer, Berlin, 1988.

- [EN] Eagon, J.A.; Northcott, D.G.: *Ideals defined by matrices and a certain complex associated with them*. Proc. R. Soc. Lond. **269**, 188-204 (1962)
- [Ei] Eisenbud, D.: *Commutative algebra with a view toward algebraic geometry*. Graduate Texts in Mathematics, Springer, Berlin, 1995.
- [EHU₁] Eisenbud, D.; Huneke, C.; Ulrich, B.: *A simple proof of some generalized principal ideal theorems*. Proc. Amer. Math. Soc. **129**, 2535–2540 (2001)
- [EHU₂] Eisenbud, D.; Huneke, C.; Ulrich, B.: *What is the Rees algebra of a module?* Proc. Amer. Math. Soc. **131**, 701–708 (2003)
- [EHU₃] Eisenbud, D.; Huneke, C.; Ulrich, B.: *Order ideals and a generalized Krull height theorem*. Math. Ann. **330**, 417–439 (2004)
- [EHU₄] Eisenbud, D.; Huneke, C.; Ulrich, B.: *Heights of ideals of minors*. Amer. J. Math. **126**, 417–438 (2004)
- [EE] Eisenbud, D.; Evans, E.G.: *A generalized principal ideal theorem*. Nagoya Math. J. **62**, 41–53 (1976)
- [FOV] Flenner, H.; O’Carroll, L.; Vogel, W.: *Joins and intersections*. Monographs in Mathematics, Springer, Berlin, 1999.
- [FL₁] Fulton, W.; Lazarsfeld, R.: *On the connectedness of degeneracy loci and special divisors*. Acta Math. **146**, 271-283 (1981)
- [Gi] Gieseker, D.: *p-ample bundles and their Chern classes*. Nagoya Math. J. **43**, 91–116 (1971)
- [EGA] Grothendieck, A.: *Éléments de géométrie algébrique*. In collaboration with J. Dieudonné, Chap. I-IV, Inst. Hautes Études Sci. Publ. Math. **4,8,11,17,20,24,28,32** (1960–1967)
- [SGA] Grothendieck, A.: *Cohomologie locale des faisceaux cohérents et théorèmes des Lefschetz locaux et globaux* (SGA2). North-Holland, Amsterdam, 1968.
- [Ha₁] Hartshorne, R.: *Ample vector bundles*. Inst. Hautes Études Sci. Publ. Math. **29**, 63–94 (1966)
- [Ha₂] Hartshorne, R.: *Algebraic geometry*. Graduate Texts in Mathematics, Springer, Berlin, 1977.
- [Jo] Józefiak, T.: *Ideals generated by minors of symmetric matrices*. Comm. Math. Helv. **53**, 595–607 (1978)
- [JP] Józefiak, T.; Pragacz, P.: *Ideals generated by Pfaffians*. J. Algebra **61**, 189–198 (1979)
- [Kl] Kleiman, S.: *Ample vector bundles on algebraic surfaces*. Proc. Amer. Math. Soc. **21**, 673–676 (1969)
- [Ma] Matsumura, H.: *Commutative ring theory*. Cambridge University Press, Cambridge, 1986.
- [St] Steffen, F.: *Connectedness theorems for determinantal schemes*. J. Algebraic Geom. **8**, 169–179 (1999)
- [Tu] Tu, L.W.: *The connectedness of degeneracy loci*. Topics in algebra (Warsaw, 1988), 235–248, Banach Center Publ. **26**, Part 2, PWN, Warsaw, 1990.

FAKULTÄT FÜR MATHEMATIK DER RUHR-UNIVERSITÄT, UNIVERSITÄTSSTR. 150, GEB. NA 2/72,
44780 BOCHUM, GERMANY

E-mail address: `hubert.flenner@rub.de`

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907-1395, USA

E-mail address: `ulrich@math.purdue.edu`